


Metcalf
Hayton
1989

INTERNATIONAL ASSOCIATION FOR
GREAT LAKES RESEARCH



J. Great Lakes Res. 15(4):654-668
Internat. Assoc. Great Lakes Res., 1989

**COMPARISON OF LEECHES AND MUSSELS AS BIOMONITORS FOR
CHLOROPHENOL POLLUTION**

Janice L. Metcalfe
*National Water Research Institute
Rivers Research Branch
P.O. Box 5050, 867 Lakeshore Road
Burlington, Ontario L7R 4A6*

Alan Hayton
*Ontario Ministry of the Environment
Water Resources Branch
P.O. Box 213, 125 Resources Road
Rexdale, Ontario M9W 5L1*

COMPARISON OF LEECHES AND MUSSELS AS BIOMONITORS FOR CHLOROPHENOL POLLUTION

Janice L. Metcalfe

*National Water Research Institute
Rivers Research Branch
P.O. Box 5050, 867 Lakeshore Road
Burlington, Ontario L7R 4A6*

Alan Hayton

*Ontario Ministry of the Environment
Water Resources Branch
P.O. Box 213, 125 Resources Road
Rexdale, Ontario M9W 5L1*

ABSTRACT. *Leeches were previously shown to have unusually high bioaccumulation capacities for chlorophenols (CPs) and have, therefore, been recommended as biomonitors for CP pollution. To determine whether leeches could survive on-site exposures in cages and to compare leeches and mussels as biomonitors for CPs, two field studies were conducted. CP residues in leeches (*Nepheleopsis obscura*) exposed for 3 weeks above and below a pulp mill complex on the Rainy River, Ontario, were compared with concentrations in resident mussels, while leeches and mussels (*Elliptio complanata*) were directly compared to exposing them simultaneously in cages near a wood preserving plant on Thunder Bay Harbour. Survival of caged leeches in the Rainy River was excellent (77-87%), provided conductivity/alkalinity was within the tolerance range of the species. Leeches as far as 100 km downstream of the mills accumulated elevated levels of CPs (21-121 ng/g vs. 4-5 ng/g pre-exposure), with proportions of the various congeners reflecting those in mill effluent. Concentrations of CPs in mussels (<0.5-3.5 ng/g) were 1-2 orders of magnitude lower than in leeches. Leeches in Thunder Bay Harbour accumulated four CPs, with concentrations of the dominant pentachlorophenol (PCP) ranging from 817-5,300 ng/g near the plant to 275-375 ng/g at 100 m, 55-85 ng/g at 300 m, and <50 ng/g at 600 m beyond the discharges. In contrast, CPs were never detected in mussels from any site. Bioassays conducted on harbor sediment using fathead minnows (*Pimephales promelas*), burrowing mayflies (*Hexagenia limbata*), and leeches demonstrated that uptake and toxicity decreased with increasing distance from the plant. At all sites, PCP residues were similar in field- and lab-exposed leeches. This study demonstrated that leeches are far superior to mussels as biomonitors for CPs, and that leeches can be successfully caged and used in routine monitoring where CP pollution is suspected.*

ADDITIONAL INDEX WORDS: *Benthos, benthic environment, invertebrates, bioaccumulation, bioassay.*

INTRODUCTION

Unionid mussels bioconcentrate a wide range of pollutants and are frequently used as biomonitors for metals (Manly and George 1977, Tessier *et al.* 1984), industrial organic chemicals (Kauss and Hamdy 1985, Pugsley *et al.* 1985), and pesticides (Miller *et al.* 1966, Bedford *et al.* 1968, Leard *et al.* 1980) in freshwater environments. One of their main advantages is their ability to survive on-site

exposures in cages for weeks or even months, which allows experimental conditions to be controlled and is of obvious benefit when studying areas not normally inhabited by mussels. There is some evidence, however, that mussels may not be suitable biomonitors for chlorophenols (CPs). Caged mussels exposed to effluent from a wood preserving plant on the Kaministikwia River in northwestern Ontario accumulated no traces of

CPs, even though concentrations were measurable in both water and caged fish (Ontario Ministry of the Environment, unpublished data).

Leeches have excellent potential as screening organisms for CPs in freshwater systems, mainly because of their unusually high biocentration capacities for these compounds. In earlier work, Metcalfe *et al.* (1984) found that leeches from an industrially polluted creek accumulated CP concentrations that were up to two orders of magnitude higher than those in fish, tadpoles, crayfish, molluscs, and a variety of aquatic insects. In a survey for the occurrence of CPs in several New Brunswick rivers, 12 chlorophenol congeners were detected in resident leeches but only three in river water (Metcalfe *et al.* 1985), indicating that leeches were the more sensitive monitoring medium. Jacob (1986) was the first to attempt caging leeches, successfully using *Haemopsis marmorata* (F. Hirudinidae) in 1-week field exposures as biomonitors for CPs in the Fraser River Estuary.

The purpose of the present study was to compare leeches and mussels as biomonitors for CP pollution originating from two industrial point sources. The distribution of CPs in the Rainy River above and below two pulp and paper mills and in Thunder Bay Harbour adjacent to a wood preserving plant was assessed using residues accumulated by caged leeches and resident mussels (Rainy River study) or caged leeches and caged mussels (Thunder Bay study). Laboratory bioassays were also conducted on contaminated sediment from the harbor to determine its toxicity and to compare CP bioaccumulation under field and laboratory conditions. Procedures for conducting field studies with caged leeches are not well established; therefore, this study contributed to the development of an appropriate protocol for these organisms.

MATERIALS AND METHODS

Study Locations

Two of the main sources of entry of CPs into aquatic environments are pulp and paper mills and wood preserving plants. In addition, CPs can be generated as a result of aqueous chlorination of organic compounds during the treatment of municipal sewage (Jones 1981). Two locations in northwestern Ontario were selected for study, both influenced by an industrial point source of CP pollution.

The first study was conducted on the Rainy

River, which flows for approximately 130 km from Rainy Lake to Lake of the Woods (Fig. 1). The main sources of CPs are two kraft pulp mills located directly across the river from each other in Fort Frances, Ontario, and International Falls, Minnesota. Minor sources are the municipal wastewater treatment plants for the towns of Fort Frances, International Falls, Emo, Rainy River, and Baudette, which serve a total population of 20,000. The mills, however, account for about 95% of the loadings of chlorinated effluents to the river in terms of discharge volume, total suspended solids and BOD (International Rainy River Water Pollution Board 1985). The focus of our second study was a wood preserving plant on Lake Superior in Thunder Bay (Fig. 2). The facility applies creosote, pentachlorophenol (PCP), and chromated copper arsenate, under pressure and heat, to lumber intended for outdoor use. Approximately 700,000 L of 5% PCP solution and 8,500,000 L of 50% creosote were used in 1983 (Bérard and Tseng 1986).

Rainy River Study

Collection, Transportation and Caging of Leeches—Approximately 600 specimens of the "bait leech," *Nepheleopsis obscura* (F. Erpobdellidae), were purchased from a bait shop in Sudbury. Their origin was natural lake populations in Wisconsin or North Dakota. Specimens were transported to the field sites in bioassay bags containing 5 L of water. The bags were given a head of oxygen, which was renewed daily, and were stored at ambient temperature. With the exception of a few injured animals, no mortality occurred during the week's field activities.

Leech cages were cylindrical in shape, 35 cm long \times 15 cm in diameter, and were constructed of 10 \times 10 mesh (10 spaces per linear inch) stainless steel wire cloth having a wire diameter of 0.064 cm and an opening of 0.191 cm. The overlapping edges were woven together with stainless steel wire, and the ends were capped with polyethylene plugs which had been pre-soaked in a cleaning solution to remove any soluble organic compounds. The plugs were secured with stainless steel gear clamps. A third clamp, with a smaller clamp attached to it, was placed around the middle of the cage. A line of quarter-inch nylon rope, strung through the smaller clamp, was used to tether the cages in the field. Our design was based on a cage described by




FIG. 1. Rainy River study sites (● = caged leeches; ○ = resident leeches; △ = resident mussels; □ = resident caddisfly larvae).




FIG. 2. Locations of leech and mussel cages in Thunder Bay Harbour (sediment bioassay sites circled).

Jacob (1986). Leeches were unable to escape from these cages in laboratory aquaria.

Experimental Design and Procedure—In early August 1986, cages containing 30 leeches (total biomass per cage approx. 90 g; average weight 2.8 g) were set at 17 sites along the river, in Rainy Lake and Lake of the Woods, and in the two major American tributaries (Fig. 1). Cages were attached by rope to either a concrete block or a navigation marker, and they rested on the substrate. Water temperature was about 18°C throughout the system. Leeches were exposed on-site for 3 weeks.

When the cages were retrieved, a dive search for resident mussels was conducted. *Lampsilis radiata siliquoidea* (Subf. Lampsilinae) were found at sites 4, 18, 28, 29, 31, and 32, while *Andonta grandis* (Subf. Anodontinae) occurred at sites 4, 31, and 32. Resident leeches, *Haemopsis grandis* (F. Hirudinidae), were also collected from sites 1 and 3 in Rainy Lake. Net-spinning caddisfly larvae of the genera *Hydropsyche* (F. Hydropsychidae) and *Neureclipsis* (F. Polycentropodidae) were found attached to the leech cages at sites 7, 8, 10, 13, 16, and 17 downstream and within 5 km of the mills. Proportionately more *Neureclipsis* were found on the cages nearest to the mills.

Samples of biota to be analyzed for CP residues were wrapped in prefired (450°C) aluminum foil, sealed in plastic bags, and frozen on dry ice at the time of collection. The following samples were prepared: duplicate composite samples of 8–11 leeches weighing 18–30 g from each site where they survived; background samples of 15 and 25 leeches weighing 38 and 82 g, respectively; composite samples of two to four specimens of each species of resident mussel, weighing 51–79 g, from each site; three resident leeches from site 1 weighing 25 g and one leech from site 3 weighing 4 g; two composite samples of caddisfly larvae from sites 7, 8, and 17 below the American mill (122 specimens weighing 7 g) and sites 10, 13, and 16 below the Canadian mill (112 specimens weighing 6 g).

Samples of river water from sites 3, 18, and 32 and the final effluents from both mills were also analyzed for dissolved and particle-bound CPs. Water or effluent was clarified on site using a stainless steel Westfalia separator. A March 5C-MD submersible magnetic drive pump was operated at a rate of 6 L/min to supply water to the centrifuge using a Teflon transfer line. One-liter samples of clarified water were acidified to pH 2.0 with sulphuric acid, then stored at 4°C until ana-

lyzed. Suspended solids were removed from the centrifuge bowls by scraping the walls with a Teflon spatula. The slurry was then pressure-filtered through a 5 µm Teflon filter and the solids transferred to a pre-cleaned glass jar where they were stored frozen until analyzed.

Analytical Methods—Biota samples were partially thawed, weighed, ground with prefired (450°C) anhydrous sodium sulphate, and Soxhlet extracted with dichloromethane for 3 hours. The extracts were evaporated at 2 mL under dry nitrogen, and lipids were removed by gel permeation chromatography using a 55/45 mixture of hexane/dichloromethane. The lipid-free sample was then extracted three times with 40, 30, and 30 mL of 0.1M potassium carbonate. To derivatize the phenols, 1 mL of acetic anhydride and 10 mL of hexane were added to the aqueous fraction and the mixture was shaken for 1 h. Finally, the hexane extracts were eluted through a micro-column of 5% deactivated silica gel with a 95/5 solution of hexane/acetone, evaporated to an appropriate volume, and analyzed for 21 CPs on a 30-m DB-5 capillary column by GC-MSD (HP5890/5970).

Suspended solids from the two pulp mill effluents were thawed and homogenized. Twenty-gram subsamples were prepared and analyzed as described for biota except that no GPC clean-up was needed, and the aqueous fraction was filtered through Celite to remove suspended material prior to acetylation.

Suspended solids from the three river water samples were prepared according to a method developed by Lee *et al.* (1987). It differed from the above-described procedure in that the 20-g subsample was first acidified to pH 1.0 with sulphuric acid, then "wet" extracted with acetone-hexane for 20 h in a Soxhlet extractor. The extract was base-partitioned with 2% potassium bicarbonate solution, and the phenols derivatized with acetic anhydride while being extracted into petroleum ether. Clean-up and analysis were as described for effluent solids. The major difference between the two methods is that the initial acidification step was omitted during preparation of the effluent solids samples. Chlorophenols tend to bind to the organic material in sediments in the undissociated form (Xie 1983). Extraction with an organic solvent will efficiently remove the CPs, provided that the pH of the sediment is acidic or neutral (M. E. Fox, National Water Research Institute, personal communication). As whole effluent from both

mills is neutral or slightly acidic (International Rainy River Water Pollution Board 1985, 1987), acidification of these samples was not considered necessary. It is possible, however, that CP extraction efficiencies were slightly higher for the river water suspended solids samples. Analytical detection limits for all of the above samples were 0.5 ng/g wet weight for all CP compounds.

Clarified water samples were analyzed for CPs according to the method of Lee *et al.* (1984). They were extracted with dichloromethane and base-partitioned with 2% potassium bicarbonate. Further preparation and analysis were as previously described for river water suspended solids samples. Detection limits were approximately 0.01 $\mu\text{g/L}$.

Thunder Bay Harbour Study

Field Exposures of Caged Leeches and Mussels—Ten sites were selected for study (Fig. 2). Sites 1, 9, and 10 were in the immediate vicinity of the western, central, and eastern discharges of the plant, respectively. Sites 8 and 16 were 100–150 m offshore, sites 7, 22, and 27 were 300–350 m offshore, site 24 was 600 m from the discharges, and the control site was approximately 3 km from the plant. Leeches (*N. obscura*) were obtained from a bait shop in Toronto which was supplied by the same wholesaler as the Sudbury bait shop where leeches were purchased for the Rainy River study. Mussels (*Elliptio complanata*, Subf. Ambleminae), aged 9–11 years and ranging from 65–72 mm in valve length, were collected from Balsam Lake in central Ontario. They were transported in plastic-lined 20-L bioassay buckets, under oxygen, at a density of 50/bucket. Leeches were transported in the same way at a density of 125/bucket. All animals were held at 4°C prior to exposure. In early June 1987, cages containing 40 leeches (total biomass per cage approx. 55 g; average weight 1.4 g) were set alongside cages of 5 mussels (total biomass per cage approx. 40 g; average weight 8 g) at each of the 10 study sites. Mussel cages were constructed of 1.25 cm mesh galvanized poultry netting and measured 30 × 45 cm; leech cages have already been described. Water temperature in the harbor was approximately 9°C. The animals were exposed on-site for 3 weeks, after which three composite samples of four leeches each and one individual mussel from each site were selected for analysis. At site 8, three additional mussels were taken. Samples of biota were wrapped in hexane-rinsed

aluminum foil, sealed in plastic bags, and stored on ice for 4–6 h prior to freezing.

Laboratory Sediment Bioassays—Sediment bioassays were conducted in October 1986. Sediment was collected by Ponar dredge (top 10 cm) from the six sites indicated in Figure 2. Unfortunately, site 1 could not be included because only creosote-soaked gravel was present. Samples were shipped to the laboratory in buckets lined with bioassay bags, and were stored at 4°C until tested.

Bioassays were conducted on fathead minnows (*Pimephales promelas*), burrowing mayflies (*Hexagenia limbata*), and leeches (*N. obscura*). The minnows were 4-month-old juveniles cultured in the laboratory from stock obtained from the USEPA Laboratory in Duluth, Minnesota. Mayflies (second-year nymphs) were collected from Georgian Bay in Lake Huron and maintained in the lab for a short period prior to testing. Leeches were purchased from the Toronto bait shop which also supplied the specimens for the field study. Bioassays were performed in 1.5-L Mason jars. A 200-mL sediment sample was placed in the jar, topped with 800 mL dechlorinated tap water, and allowed to settle for 24 h. The system was then aerated and the test organisms introduced. Tests were conducted separately and in triplicate for each organism, using five minnows, five mayflies, or three leeches per jar. Due to a shortage of specimens, tests on sediment from sites 9 and 10 could not be replicated for mayflies or leeches. Bioassays were run for 10 days. Mortality was recorded daily and dead organisms were removed and discarded. After 10 days' exposure, surviving animals from each treatment (site/species) were composited, wrapped in hexane-rinsed aluminum foil, sealed in plastic bags, and stored frozen until analyzed for CPs.

Analytical Methods—Samples of field-exposed biota were weighed and homogenized using a sonification process. The samples were then digested with HCl and extracted with dichloromethane. Lipids were removed and the lipid-free extracts were dried through sodium sulphate and concentrated on a rotary evaporator. Extracts were derivatized using diazomethane, subjected to a Florisil clean-up, concentrated on a vortex evaporator, and analyzed for six CPs using a capillary column gas chromatograph with an electron capture detector. Detection limits for this method were 100 ng/g

wet weight for 2,3,4-TCP (TCP = trichlorophenol) and 50 ng/g for all other compounds.

Samples of laboratory-exposed biota were weighed and homogenized. An aliquot was digested overnight by gentle agitation with hydrochloric acid. The sample was then extracted with a 25/75 mixture of dichloromethane/hexane. The extract was transferred to a gel permeation column and eluted with a solution of 45/55 dichloromethane/hexane to remove the lipids. The lipid-free extract was concentrated, derivatized with diazomethane and subjected to a Florisil clean-up. The resulting extract was evaporated to an appropriate volume and analyzed for six CPs by dual capillary column/dual ECD gas chromatography. Method detection limits were 5 ng/g wet weight for 2,3,4,5-TECP (TECP = tetrachlorophenol) and PCP, and 10 ng/g for the remaining compounds.

RESULTS

Rainy River Study

At 11 of the 17 caged leech sites, no leeches survived the 3-week exposure period. At the three sites above the mills (1, 4, and 5), leeches had apparently been dead for some time, as only traces of tissue were left in the cages. At the six sites downstream and within 5 km of the mills (7, 8, 10, 13, 16, and 17), all leeches were dead and covered with fungus but had not completely decomposed. At sites 26 and 28 (mid-river), all leeches were dead, but there was no evidence of fungus or decomposition, which suggests they had been dead only a few days. Survival at the remaining six sites was excellent, ranging from 77–87%. These sites included four in the lower river (33, 34, 31, and 32), one in the Big Fork River (24), and one in the Baudette River (30).

Prior to setting the cages in the river, it was noted that many of the leeches were clitellate, that is, in reproductive condition. During the exposure period, many leeches deposited cocoons on the end plugs of their cages. Numerous cocoons were seen at sites 5, 7, 8, 10, 13, 26, 31, and 33 with lesser numbers at the remaining sites except for sites 1 and 4 where none were found.

Concentrations of CPs in clarified water and suspended solids from sites 3, 18 and 32 and both final mill effluents are presented in Table 1. The Canadian mill effluent contained much higher concentrations of total CPs (6.09 $\mu\text{g/L}$ in water; 353.9 ng/g in suspended solids) than the American mill effluent

(0.25 $\mu\text{g/L}$ in water; 5.8 ng/g in solids). Dominant congeners in the dissolved and particle-bound phases of both effluents were 2,4,6-TCP, 2,3,4,6-TECP, and PCP, accounting for 73–100% of the total CPs. Suspended solids from the Canadian effluent also contained 19% 2,4-DCP (DCP = dichlorophenol). Traces of 4-MCP (MCP = monochlorophenol) and 2,4,6-TCP in clarified water and PCP in suspended solids were found in the river both upstream and 5 km downstream of the mills, whereas only minute traces of 4-MCP were found in water from Lake of the Woods.

Figure 3 compares the concentrations of total CPs accumulated by caged and resident leeches with concentrations in resident mussels. No data are shown for caddisfly larvae because they did not contain detectable residues, even though they were collected from sites located within 5 km of the mills. Concentrations of CPs in mussels were very low, ranging from <0.5 to 3.5 ng/g. The highest value was for *L.r. siliquoidea* collected from site 18, 5 km below the mills. Both species of mussels were collected from sites 4, 31, and 32, but the *L.r. siliquoidea* specimens from sites 4 and 32 were lost in a freezer malfunction; therefore, interspecific comparisons were not possible. Concentrations of total CPs in leeches prior to exposure were approximately 5 ng/g, which is similar to concentrations in resident leeches from Rainy Lake. After 3 weeks' exposure, concentrations in leeches increased to 21–121 ng/g at the four sites in the lower river, but *decreased* to less than 2 ng/g in the Big Fork River and to about 3 ng/g in the Baudette River.

Figure 4 compares relative proportions of the various CPs in leeches with proportions in effluent from the major point source—the Canadian mill. The pattern for the American mill effluent was very similar. For clarity, only congeners accounting for over 5% of the total CPs are included. The average profiles for caged leeches from the four sites in the lower Rainy River closely resembled the pattern for mill effluents, while those for caged leeches from the two American tributaries did not. Rainy Lake residents contained mainly PCP and some 2,3,4,6-TECP. The pattern in pre-exposure leeches was unique, with 4-MCP, 2,4-DCP, 2,4,6-TCP, and PCP present in almost equal portions. In Figure 5, proportions of CPs in resident mussels are compared with the effluent profile. The pattern for *L.r. siliquoidea* collected farther downstream contained proportionately more of the lower chlorinated compounds (mono- and dichlorophenols), as did *A. grandis* from all sites. Pentachlorophenol was never

TABLE 1. Concentrations of chlorophenols in clarified water ($\mu\text{g/L}$) and suspended solids (ng/g wet weight) from mill effluents and river water samples (no value = not detectable).

Chlorophenol isomer	Canadian mill effluent		American mill effluent		Site 3		Site 18		Site 32	
	Water	Solids	Water	Solids	Water	Solids	Water	Solids	Water	Solids
2-MCP	0.015	1.3								
4-MCP	0.014	1.2	0.011		0.011		0.011		0.006	
2,6-DCP	0.056	8.8								
3METHYL-4CP		1.5								
2,4-DCP	0.162	65.8	0.021							
3,5-DCP	0.015	2.9								
3,4-DCP		1.6								
2,4,6-TCP	4.299	208.0	0.145	2.5	0.030		0.050			
2,4,5-TCP		3.6								
2,3,4-TCP		2.0								
3,4,5-TCP		3.9								
2,3,4,6-TECP	1.064	39.7	0.041	1.8						
2,3,4,5-TECP		1.3								
PCP	0.465	12.3	0.028	1.5		0.9		1.1		
TOTAL	6.090	353.9	0.246	5.8	0.044	0.9	0.061	1.1	0.006	




FIG. 3. Concentrations of total chlorophenols (ng/g wet weight) in leeches (numbers in lower portion of figure) and mussels (numbers in upper portion) in the Rainy River (values for caged leeches represent duplicate samples).




FIG. 4. Relative proportions of chlorophenol congeners in leeches and in pulp mill effluent. (For effluent, dark and light bars represent suspended solids and clarified water, respectively.)

detected in mussels, yet this compound was an important congener in caged and resident leeches.

Thunder Bay Harbour Study

All caged mussels survived the 3-week exposures in Thunder Bay Harbour, but with one exception did not accumulate detectable concentrations of any CP compound at any site. The exception was a mussel from site 8, which contained 60 ng/g 2,4,5-TCP, 85 ng/g 2,3,4,5-TECP, and 930 ng/g PCP. Three additional mussels from this site were analyzed individually and found to contain no traces of CPs. We therefore suspect that the original sample had been contaminated or mislabelled during preparation.

Caged leeches suffered some mortality at all sites, not exceeding 38% except at site 10 where 45% were dead and the rest moribund. It was also observed that specimens at site 1 lacked vigor. Unlike mussels, leeches did bioaccumulate CPs (Fig. 6). In decreasing order of importance, PCP, 2,3,4,5-TECP, 2,4,5-TCP, and 2,4,6-TCP were identified in leech tissues. Concentrations of PCP ranged




FIG. 5. Relative proportions of chlorophenol congeners in mussels and in pulp mill effluent. (Legend as per Fig. 4.)

from 817 to 5,300 ng/g at sites 1, 9, and 10, decreased to 275-375 at sites 8 and 16, decreased further to 55-85 ng/g at sites 7, 22, and 27, and were undetectable (<50 ng/g) at sites 24 and the control. Concentrations of 2,3,4,5-TECP and 2,4,5-TCP were 63-427 ng/g and 57-90 ng/g, respectively, in leeches caged near the discharges. Unlike PCP, however, these compounds did not persist into the harbor. A different distribution pattern was observed for 2,4,6-TCP. Concentrations of this isomer were similar at all sites (<50-70 ng/g) including the control. Leeches did not accumulate 2,3,4-TCP or 2,3,5,6-TECP at any site. All values reported above are averages for three composite samples of leeches. Variability was low, with stan-




FIG. 6. Concentrations of four chlorophenol congeners (ng/g wet weight) in caged leeches from Thunder Bay Harbour.

dard deviation generally 10–20% of the mean. For the purpose of calculating averages, all values < 50 ng/g were considered to be equal to 50 ng/g.

All minnows, mayflies, and leeches survived 10-d exposures in the laboratory to sediment from sites 7, 8, and the control. Two of 15 mayflies and one of nine leeches died after exposure to sediment from site 24, and all but one organism died after exposure to sediment from sites 9 and 10. The survivor was a leech from site 9. Concentrations of tri- and tetrachlorophenols in the tissues of surviving animals were near or below the detection limit for minnows and mayflies from all sites (Table 2). In leeches, concentrations of 2,4,5-TCP and the tetrachlorophenols were elevated only at site 9, the central discharge, while concentrations of 2,4,6-TCP were similar at all sites. Concentrations of

PCP in leeches dropped off in a transect from the central discharge into the harbor. A similar trend was suggested by the minnow and mayfly data. Wherever data for all three test organisms were available, i.e., sites 8, 7, 24, and the control, leeches were shown to accumulate the highest concentrations of 2,4,6-TCP and PCP.

DISCUSSION

The results of the Rainy River study demonstrated that field studies using caged leeches are feasible. Excellent survival rates were observed at six of the 17 sites, which indicates that the handling procedures and cage design were suitable and the exposure period appropriate. The cause of high mortality at the remaining 11 sites is believed to be related

TABLE 2. Concentrations of chlorophenols (ng/g wet weight) in organisms surviving the sediment bioassays.

Site	Leeches					
	2,4,6-TCP	2,4,5-TCP	2,3,4-TCP	2,3,5,6-TECP	2,3,4,5-TECP	PCP
9	120	5,780	20	220	843	1,080
8	140	<10	20	30	<5	102
7	120	60	<10	50	6	92
24	110	30	20	30	8	42
Control	180	50	50	40	13	79
	Mayflies					
9	-*	-	-	-	-	-
8	30	30	<10	20	<5	68
7	40	30	<10	20	<5	31
24	30	50	20	30	8	66
Control	30	10	10	20	6	20
	Minnows					
9	-	-	-	-	-	-
8	20	<10	<10	10	<5	61
7	<10	10	<10	<10	<5	13
24	30	20	<10	20	<5	42
Control	30	30	<10	10	9	23

*dead, no sample

to water quality, and this will be discussed in detail below.

The Canadian pulp mill is obviously the major source of CPs to the Rainy River. Concentrations of total CPs in clarified water and suspended solids were 25× and 60× higher, respectively, than in effluent from the American mill. Assuming these samples are representative, actual daily contributions to the river would be about 430 g total CPs for the Canadian mill and only 30 g for the American mill, based on discharge volumes of 70 × 10⁶L/d and 124 × 10⁶L/d and suspended solids loadings of 9,300 kg/d and 3,272 kg/d, respectively, in 1986 (International Rainy River Water Pollution Board 1987). Concentrations of CPs in river water and suspended solids both upstream and 5 km downstream of the mills were similar, suggesting that the impact zone of the pulp mill complex extended no farther than 5 km.

Concentrations of CPs in caged leeches increased above background levels in animals exposed in the lower Rainy River, but decreased in those exposed in the Big Fork and Baudette rivers. Although the latter site was just below the town of Baudette's sewage treatment plant, concentrations in leeches were very low. Site 34 in the Rainy River was also below a sewage treatment plant (for the

town of Rainy River), yet concentrations in leeches here were no higher than in those upstream of the plant. For these reasons, local municipal discharges did not appear to be responsible for the elevated levels of CPs in leeches in the lower Rainy River. Rather, this contamination is most likely attributed to the pulp mills located over 100 km upstream. Resident mussels collected throughout the Rainy River accumulated CP concentrations that were 1–2 orders of magnitude lower than those in caged leeches from the lower river. Mussels have been previously shown to have lower bioconcentration factors (BCFs) for CPs than leeches. Metcalfe *et al.* (1988) reported BCFs of 1,200–4,200 for 2,4,6-TCP, 4,000–5,500 for 2,3,4,6-TECP, and 3,000–4,300 for PCP in leeches from a contaminated creek, whereas BCFs reported in the literature for bivalves exposed to similar levels of contamination were 60 and 170 for 2,3,4,6-TECP and PCP, respectively, in the marine mussel *Mytilus edulis* (Folke *et al.* 1983) and 64 for 2,4,6-TCP in the freshwater mussel *Andonta piscinalis* (Paasivirta *et al.* 1985). Chlorophenols are eliminated rapidly by most aquatic organisms, including mussels. A half life of 2–3 d has been reported for PCP in the marine mussel, *M. edulis* (Ernst 1979). In contrast, half lives for these com-

pounds in leeches are generally longer than 1 month (Metcalf *et al.* 1988). Assuming a 30-d half life for total CPs in leeches, and given a 4–5 ng/g background concentration in pre-exposure leeches, the 2–3 ng/g concentrations observed in leeches from the Big Fork and Baudette rivers are consistent with residues expected after 3–4 weeks' depuration in clean water. While the elimination of CPs by leeches is very slow, uptake is known to be rapid. Jacob (1986) exposed the leech *H. marmorata* to 10 µg/L concentrations of 5 CPs in the laboratory for 1 week and observed that equilibrium conditions were achieved within 4–5 d at temperatures of 4 and 12°C, although not within 7 d at 22°C. This combination of rapid uptake and slow elimination is probably due to the absence in these primitive organisms of enzyme systems capable of conjugating CPs and will, of course, result in very high BCFs for these compounds in leeches.

Relative proportions of the various CP isomers in pre-exposure leeches were dramatically altered after 3-week exposures in the field, presumably in response to local conditions. For example, profiles in caged leeches from all sites in the lower Rainy River closely reflected the pattern for effluent from the Canadian mill, which is the major source of CPs to the river. Profiles for caged leeches from the American tributaries and resident leeches from Rainy Lake, however, were more characteristic of locations remote from point sources of CPs. These specimens contained high proportions of PCP, which is ubiquitous in the environment (Jones 1981), and which was shown to be the major isomer in suspended solids from Rainy Lake. In contrast to leeches, profiles for mussels from Rainy Lake were essentially the same as those for mussels collected downstream of the mills. While leeches accumulated mainly 2,4,6-TCP, 2,3,4,6-TECP, and PCP, mussels contained mainly di- and trichlorophenols, very little 2,3,4,6-TECP, and no PCP. Paasivirta *et al.* (1985), in their study on a CP-contaminated lake chain in Finland, also observed that mussels and other filter-feeding benthic invertebrates selectively accumulated the more water soluble compounds such as 2,4-DCP and 3,4-DCP.

The pattern of mortality observed among caged leeches in the Rainy River does not implicate the pulp mills as the cause, because 100% mortality occurred at all sites above the mills and 50–80 km downstream in addition to the six sites within 5 km of the outfalls. Instead, it suggests that differences in water quality between the upper and lower reaches of the river and between the main river and

its tributaries are more likely to be responsible. To determine which factors may be involved, water quality data for the study area (Table 3) were compared with the environmental requirements of the test species from the literature.

N. obscura is a cold-water stenotherm occurring at temperatures of 4–15°C in Alberta (Linton *et al.* 1983a), below 16°C in Michigan (Kopinski 1972), and below 24°C in Colorado (Herrmann 1970a). It is tolerant of pH fluctuations between 6.3–9.4 in Colorado (Herrmann 1970a) and 7.0–7.5 in Michigan (Klemm 1972). Herrmann (1970b) reported a tolerance range of 11.8–654.4 mg/L total dissolved solids (TDS) for *N. obscura*, which is both narrow and low. In laboratory experiments, Reynoldson and Davies (1980) observed optimal survival of this species at conductivities ranging from 1,584–5,554 µmhos/cm, and high mortality below 84 µmhos/cm. Most leech species are most abundant at alkalinity above 60 mg/L, while few are present below 18 mg/L (Sawyer 1974). Water quality conditions in the Rainy River were apparently suboptimal for *N. obscura*. Temperatures throughout the system were near the upper tolerance limit for the species. Conductivity, alkalinity, and TDS were at or below its lower tolerance limit in the upper river, although conditions were more favorable in the lower river and the tributaries. Although these differences appear minor, they may have been critical considering the life stage of the test specimens. In Minnesota, *N. obscura* has a 24-month life cycle (Peterson 1983). Clitellum development requires a body weight of 1.20 g, and mature leeches die shortly after depositing cocoons. They are clitellate in June–July and exhibit a lack of normal vigor and poor body condition by July–August when reproduction is complete or nearing completion. Test specimens, which were obtained from lakes in Wisconsin or North Dakota in early August, were in reproductive condition (clitellate; body weight of 1.5–4.0 g). Several studies have reported narrower tolerance ranges for both temperature (Linton *et al.* 1983a) and dissolved salts (Reynoldson and Davies 1980) in post-reproductive individuals. It is therefore hypothesized that because the animals were already under stress due to high water temperatures and their reproductive activities, the slightly lower values for conductivity, alkalinity, and TDS in the upper Rainy River were responsible for the high mortality observed at these sites.

Cocoon production by caged leeches at various sites demonstrated an interesting phenomenon. Although environmental conditions were less than

TABLE 3. Rainy River water quality data for August, 1986 (range for summer months in brackets).

Location	% survival of caged leeches	Temp (°C)	pH	Conductivity (µmhos/cm)	TDS (mg/L)	Alkalinity (mg/L CaCO ₃)
Rainy Lake	0	21.0	7.2	- (47- 51)	-	-
Above mills	0	21.0	7.0	49 (47- 51)	39 (39-73)	19 (19)
15 km below mills	N*	22.5	7.7	68 (54- 68)	52 (50-75)	24 (18-24)
Above Emo	N*	22.0	6.9	72 (61- 72)	36 (36-65)	25 (22-30)
Below Emo	0	21.5	7.0	72 (61- 72)	50 (50-70)	25 (22-28)
Above Rainy River	83	23.0	6.9	84 (71- 84)	62 (62-80)	31 (29-33)
Below Rainy River	87	22.0	6.9	84 (71- 84)	62 (62-85)	31 (28-31)
Big Fork River	87	23.0	8.0	- (140-210)	-	-
Baudette River	80	22.0	7.3	- (120-180)	-	-

*N = no cage set

No value = no data available

ideal, cocoons were deposited—frequently in large numbers—at most sites throughout the river. Linton *et al.* (1983b) demonstrated that reproduction, once it has begun, will continue even under suboptimal conditions because the adults have a limited life span once committed to this activity. As the test leeches were apparently well into this phase of their life cycle when obtained from the bait shop, this explains why cocoons were found at most sites. The reason why no cocoons were found at sites 1 and 4 may be that conditions were almost immediately lethal to the adults.

Several recommendations for conducting successful field studies with caged leeches, particularly *N. obscura*, arose from the Rainy River study. First of all, specimens in reproductive condition should be avoided, as they are nearing the end of their life cycle and natural mortality rates will be high. Furthermore, they are more sensitive to suboptimal environmental conditions during this phase. Secondly, studies should be conducted at a time of year when water temperatures are cooler and, therefore, more favorable. Finally, the concentration of dissolved salts in the water is a primary limiting factor for leeches (Herrmann 1970a) and should be given prior consideration. Applying what was learned to the study on Thunder Bay Harbour, smaller, prereproductive leeches were used and the study was conducted in early June when water temperature (9°C) was more acceptable to the test species. Conductivity and alkalinity throughout the harbor were also above the minimum requirements for *N. obscura*, at about 110 µmhos/cm and 45 mg/L, respectively.

In the Rainy River study, the biomonitoring potential of caged leeches was compared with that of resident mussels. The Thunder Bay Harbour field study provided a more direct comparison of these two invertebrates, because both organisms were caged and exposed under identical field conditions. Nevertheless, the findings of both studies were similar. Caged mussels did not bioaccumulate CPs above detection limits at any site in Thunder Bay Harbour; therefore, residues in mussels provided no information on the distribution of CPs in the receiving waters adjacent to the wood preserving plant. Leeches, however, accumulated four CP congeners in concentrations ranging from <50 to 5,300 ng/g. From the patterns of residues in leech tissues, the extent of contamination originating from the plant could be determined, the plant was implicated as the source of 2,4,5-TCP, 2,3,5,6-TECP, 2,3,4,5-TECP, and PCP (but not 2,4,6-TCP) to the harbor, PCP was identified as the major and most persistent congener, and the relative loadings of the three plant discharges could be distinguished from one another.

Leeches were apparently more sensitive than mussels to contaminants discharged from the plant, as they suffered some mortality at all field sites. Field conditions may have been more stressful to leeches than lab conditions, however, as mortality in the lab occurred only after exposure to sediment from the most contaminated sites. Leeches exposed to sediment from site 9 in the lab accumulated higher concentrations of 2,4,6-TCP, 2,4,5-TCP, 2,3,4,5-TECP, and 2,3,5,6-TECP than leeches exposed in the field, i.e., 120 vs. 57 ng/g,

TABLE 4. Bioaccumulation of pentachlorophenol (ng/g wet weight) from Thunder Bay Harbour sediment under field and laboratory conditions (no value = not tested).

Site	Field	Laboratory		
	Caged Leeches	Leeches	Mayflies	Minnows
1	5,300	--	--	--
10	1,267	DEAD	DEAD	DEAD
9	817	1,080	DEAD	DEAD
8	373	102	68	61
7	83	92	31	13
24	<50	42	66	42
Control	<50	79	20	23

5780 vs. 57 ng/g, 843 vs. 63 ng/g, and 220 vs. <50 ng/g, respectively, although concentrations of PCP were similar (1,080 vs. 817 ng/g). This suggests that leeches were in closer contact with contaminated sediment in the lab. In most other aspects, results of the field and laboratory studies correlated well. Concentrations of 2,4,6-TCP accumulated by leeches, mayflies, and minnows in the lab and by caged leeches in the field showed no site-to-site variations, implying that this compound is ubiquitous in the area and is not associated with the plant's discharges. In leeches, concentrations of 2,4,5-TCP, 2,3,4,5-TECP, and 2,3,5,6-TECP were elevated only in specimens from sites 1, 9, and 10 in the field and site 9 in the laboratory, which indicates that these contaminants do not persist beyond the immediate vicinity of the discharges. A comparison of PCP bioaccumulation under field and lab conditions is presented in Table 4. At all sites, PCP residues were similar in both field- and lab-exposed leeches. The data for all organisms showed a trend toward decreasing uptake and decreasing toxicity in a transect away from the discharges. This does not necessarily identify PCP or CPs in general as the cause of toxicity. The analysis of selected organic and inorganic toxicants in sediments near the plant revealed high concentrations of CPs (approximately 5,000 ng/g dry weight), chlorobenzenes (90 ng/g), DDE (25 ng/g), octachlorostyrene (15 ng/g), As (25 µg/g), Cr (100 µg/g), Cu (80 µg/g), and Zn (350 µg/g) (Ontario Ministry of the Environment, Northwestern Region, unpublished data). The Ontario Ministry of the Environment's criteria for open-water disposal of dredged materials (Thomas and Mudroch 1978) were exceeded for the four metals. Sediments are also highly contam-

inated with oil and grease. Samples collected in 1984 contained 7,600–80,000 µg/g oil and grease (Bérard and Tseng 1986), greatly exceeding OME's dredging guidelines of 1,500 µg/g.

CONCLUSIONS

These studies on the Rainy River and Thunder Bay Harbour have shown that environmental monitoring with caged leeches is feasible, provided the life history of the test species is considered and its specific environmental requirements are accommodated. Leeches in reproductive condition should be avoided, as their poor condition and lowered resistance will lead to high mortality. Concentrations of dissolved salts and temperature appear to be critical water quality factors affecting survival. Incompatibility between test organism and study area could be avoided by recruiting local species, if they are available in sufficient numbers.

Caged leeches exhibited some mortality after exposure in Thunder Bay Harbour, while mussels subjected to identical conditions did not. Leeches may simply be less hardy than mussels, which would be a drawback to their use in field studies, or they may have been more sensitive to the specific toxicants discharged from the wood preserving plant.

Leeches were shown to have much higher bioconcentration capacities for CPs than resident mussels in the Rainy River and caged mussels in Thunder Bay Harbour. There was also evidence that leeches accumulate higher concentrations of CPs than net-spinning caddisfly larvae, burrowing mayflies, and fathead minnows. Leeches may therefore be the best available screening organism for CPs in aquatic environments.

Over a 3-week exposure period in the field, leeches appeared capable of "adjusting" their body burdens to reflect the concentrations and relative proportions of CPs in the environment to which they were introduced. Exposures of shorter duration may be adequate, however, as 10 days were sufficient to demonstrate site-to-site differences in CP uptake from contaminated sediment and, according to Jacob (1986), equilibrium with waterborne CPs may be reached within a week. Because leeches eliminate CPs very slowly, any residues present in their tissues initially could persist and lead to erroneously high final concentrations. Screening the stock of test organisms for CPs prior to exposure, to be sure they are contaminant-free, is therefore strongly recommended.

The origin, distribution, and persistence (with respect to bioavailability) of CPs in Thunder Bay Harbour could be determined from residues of CPs accumulated by caged leeches. Results of laboratory sediment bioassays supported the main findings of the field study, and also demonstrated that toxicity increased with increasing sediment contamination. When used together, bioaccumulation and toxicity studies can provide a powerful bioassessment tool for determining the impact of point sources of pollution on their receiving waters.

ACKNOWLEDGMENTS

The authors wish to thank D. Hollinger and J. Vander Wal, Ontario Ministry of the Environment, Northwestern Region, for excellent logistics support including the use of their vehicles and vessels for the Rainy River study. Rainy River biota samples were prepared by S. P. Batchelor and C. Ferguson of the National Water Research Institute, and analyzed by A. Lumb and E. Chisholm of the Water Quality National Laboratory in Burlington, Ontario. Data on CPs in Rainy River water and suspended solids were provided by John C. Merriman, Water Quality Branch, Ontario Region. The Ontario Ministry of the Environment, Northwestern Region, and the Minnesota Pollution Control Agency kindly provided us with water quality data for the Rainy River. Field support for the Thunder Bay Harbour study was provided by B. Shadford, and biota samples were analyzed by the Laboratory Services Branch, Ontario Ministry of the Environment, and by Enviroclean Division of MacLaren Plansearch Inc.

REFERENCES

- Bedford, J. W., Roelofs, E. W., and Zabik, M. J. 1968. The freshwater mussel as a biological monitor of pesticide concentrations in a lotic environment. *Limn. Oceanogr.* 13:118-126.
- Bérard, M.-F., and Tseng, T. 1986. *Survey of Northern Wood Preservers Inc., Thunder Bay, Ontario, October 1984*. Environment Canada, Environmental Protection, Pollution Abatement Division, Toronto, Ontario.
- Ernst, W. 1979. Factors affecting the evaluation of chemicals in laboratory experiments using marine organisms. *Ecotoxicol. Environm. Safety* 3:90-98.
- Folke, J., Birklund, J., Sorensen, A. K., and Lund, U. 1983. The impact on the ecology of polychlorinated phenols and other organics dumped at the bank of a small marine inlet. *Chemosphere* 12:1169-1181.
- Herrmann, S. J. 1970a. Systematics, distribution, and ecology of Colorado Hirudinea. *Amer. Midl. Natur.* 83:1-37.
- _____. 1970b. Total residue tolerances of Colorado Hirudinea. *Southwest. Natur.* 15:269-272.
- International Rainy River Water Pollution Board. 1985. 33rd progress report, IRRWPB, International Joint Commission.
- _____. 1987. 35th progress report, IRRWPB, International Joint Commission.
- Jacob, C. 1986. Use of the bioconcentration capability of leeches to evaluate chlorophenol pollution. M.Sc. dissertation, The University of British Columbia, Vancouver, British Columbia.
- Jones, P. A. 1981. *Chlorophenols and their impurities in the Canadian environment*. Environment Canada, Environmental Protection Service, Economic and Technical Review Report Number EPS 3-EC-81-2, Ottawa, Canada.
- Kauss, P. B., and Hamdy, Y. S. 1985. Biological monitoring of organochlorine contaminants in the St. Clair and Detroit rivers using introduced clams, *Elliptio complanatus*. *J. Great Lakes Res.* 11: 247-263.
- Klemm, D. J. 1972. The leeches (Annelida: Hirudinea) of Michigan. *Mich. Academician* 4:405-444.
- Kopinski, M. L. 1972. Leeches (Hirudinea) of Marquette County, Michigan. *Mich. Academician* 4: 377-383.
- Leard, R. L., Grantham, B. J., and Pessoney, G. F. 1980. Use of selected freshwater bivalves for monitoring organochlorines pesticide residues in major Mississippi stream systems, 1972-73. *Pestic. Monit. J.* 14:47-52.
- Lee, H. B., Weng, L. D., and Chau, A. S. Y. 1984. Chemical derivatization analysis of pesticide residues. VIII. Analysis of 15 chlorophenols in natural water by in situ acetylation. *J. Assoc. Off. Anal. Chem.* 67:789-794.
- _____, Stokker, Y. D., and Chau, A. S. Y. 1987. Analysis of phenols by chemical derivatization. V. Determination of pentachlorophenol and 19 other chlorinated phenols in sediments. *J. Assoc. Off. Anal. Chem.* 70:1003-1008.
- Linton, L. R., Davies, R. W., and Wrona, F. J. 1983a. The effects of water temperature, ionic content and total dissolved solids on *Nepheleopsis obscura* and *Erpobdella punctata* (Hirudinoidea, Erpobdellidae). 1. Mortality. *Holarct. Ecol.* 6:59-63.
- _____, Davies, R. W., and Wrona, F. J. 1983b. The effects of temperature, ionic content and total dissolved solids on *Nepheleopsis obscura* and *Erpobdella punctata* (Hirudinoidea, Erpobdellidae). 2. Reproduction. *Holarct. Ecol.* 6:64-68.
- Manly, R., and George, W. O. 1977. The occurrence of some heavy metals in populations of the freshwater

- mussel *Anodonta anatina* (L.) from the River Thames. *Environ. Pollut.* 14:139-154.
- Metcalf, J. L., Fox, M. E., and Carey, J. H. 1984. Aquatic leeches (Hirudinea) as bioindicators of organic chemical contaminants in freshwater ecosystems. *Chemosphere* 13:143-150.
- _____, Fox, M. E., Coletta, P. A., and Carey, J. H. 1985. *Chlorophenol levels in aquatic leeches from selected sites in southwestern New Brunswick*. Environment Canada, Inland Waters Directorate, National Water Research Institute, Contribution No. 85-61, Burlington, Ontario.
- _____, Fox, M. E., and Carey, J. H. 1988. *Freshwater leeches (Hirudinea) as a screening tool for detecting organic contaminants in the environment*. Environment Canada, Inland Waters Directorate, National Water Research Institute, Contribution No. 88-13, Burlington, Ontario.
- Miller, C. W., Zuckerman, B. M., and Charig, A. J. 1966. Water translocation of Diazinon-C¹⁴ and Parathion-S³⁵ off a model cranberry bog and subsequent occurrence in fish and mussels. *Trans. Amer. Fish. Soc.* 95:345-349.
- Paasivirta, J., Heinola, K., Humppi, T., Karjalainen, A., Knuutinen, J., Mantykoski, K., Paukku, R., Piilola, T., Surma-Aho, K., Tarhanen, J., Welling, L., Vihonen, H., and Sarkka, J. 1985. Polychlorinated phenols, guaiacols and catechols in environment. *Chemosphere* 14:469-491.
- Peterson, D. L. 1983. Life cycle and reproduction of *Nepheleopsis obscura* Verrill (Hirudinea: Erpobdellidae) in permanent ponds of northwestern Minnesota. *Freshwat. Invertebr. Biol.* 2:165-172.
- Pugsley, C. W., Hebert, P. D. N., Wood, G. W., Brotea, G., and Obal, T. W. 1985. Distribution of contaminants in clams and sediments from the Huron-Erie corridor. I. PCBs and octachlorostyrene. *J. Great Lakes Res.* 11:275-289.
- Reynoldson, T. B., and Davies, R. W. 1980. A comparative study of weight regulation in *Nepheleopsis obscura* and *Erpobdella punctata* (Hirudinoidea). *Comp. Biochem. Physiol.* 66A:711-714.
- Sawyer, R. T. 1974. Leeches (Annelida: Hirudinea). In *Pollution Ecology of Freshwater Invertebrates*, ed. C. W. Hart and S. L. H. Fuller, pp. 81-142. New York: Academic Press.
- Tessier, A., Campbell, P. G. C., Auclair, J. C., and Bisson, M. 1984. Relationships between the partitioning of trace metals in sediments and their accumulation in the tissues of the freshwater mollusc *Elliptio complanata* in a mining area. *Can. J. Fish. Aquat. Sci.* 41:1463-1472.
- Thomas, R. L., and Mudroch, A. 1978. *Small craft harbours—sediment survey. Lakes Ontario, Erie and Lake St. Clair*. Report to Small Craft Harbours Ontario Region from the Great Lakes Biolimnology Laboratory, Burlington, Ontario.
- Xie, T. M. 1983. Determination of trace amounts of chlorophenols and chloroguaiacols in sediment. *Chemosphere* 12:1183-1191.